



(2017). Impact of age and antibody type on progression from single to multiple autoantibodies in type 1 diabetes relatives. *Journal of Clinical Endocrinology and Metabolism*, 102(8), 2881-2886.
<https://doi.org/10.1210/jc.2017-00569>

Peer reviewed version

Link to published version (if available):
[10.1210/jc.2017-00569](https://doi.org/10.1210/jc.2017-00569)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via OUP at <https://academic.oup.com/jcem/article-lookup/doi/10.1210/jc.2017-00569#94753474>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

The Journal of Clinical Endocrinology & Metabolism

Impact of age and antibody type on progression from single to multiple autoantibodies in type 1 diabetes relatives

--Manuscript Draft--

Manuscript Number:	jc.2017-00569R2
Article Type:	Clinical Research Article
Full Title:	Impact of age and antibody type on progression from single to multiple autoantibodies in type 1 diabetes relatives
Short Title:	Autoantibody progression in pre-type 1 diabetes
Corresponding Author:	Emanuele Bosi, M.D. Universita Vita Salute San Raffaele Milan, ITALY
Corresponding Author's Institution:	Universita Vita Salute San Raffaele
Order of Authors:	Emanuele Bosi, M.D. David C Boulware Dorothy J Becker Jane H Buckner Susan Geyer Peter A Gottlieb Courtney Henderson Amanda Kinderman Jay M Sosenko Andrea K Steck Polly J Bingley
Section/Category:	Diabetes, Pancreatic and Gastrointestinal Hormones
Manuscript Classifications:	Diabetes; Pre-diabetes; Type 1 Diabetes
Keywords:	type 1 diabetes; autoantibodies; prediction; prevention.
Abstract:	<p>Context</p> <p>Islet autoantibodies are markers of type 1 diabetes and an increase in number of autoantibodies detected during the preclinical phase is predictive of progression to overt disease.</p> <p>Objective</p> <p>To refine the impact of age in relation to islet antibody type on the progression from single to multiple autoantibodies in relatives of people with type 1 diabetes.</p> <p>Research design and methods</p> <p>We examined 994 relatives with normal glucose tolerance and positive for a single autoantibody, followed prospectively in the TrialNet Pathway to Prevention. Antibodies to GAD (GADA), insulin (IAA), IA-2 (IA-2A), zinc transporter 8 (ZnT8A) and ICA were tested every 6-12 months. The primary outcome was confirmed development of multiple autoantibodies. Age was categorized as <8yr; 8-11yr; 12-17yr; ≥18yr and optimal age breakpoints identified by recursive partitioning analysis.</p> <p>Results</p>

	<p>After median follow-up of 2 years, 141 relatives had developed ≥ 1 additional autoantibodies. Five-year risk was inversely related to age, but the pattern differed by antibody type: relatives with GADA showed a gradual decrease in risk over the four age groups, while relatives with IAA showed a sharp decrease above the age of 8 years. Recursive partitioning analysis identified age breakpoints at 14 years in relatives with GADA and at 4 years in relatives with IAA.</p> <p>Conclusions</p> <p>In relatives with IAA, spread of islet autoimmunity is largely limited to early childhood, while immune responses initially directed at GAD can mature over a longer period of time. These differences have important implications for monitoring these subjects and for designing prevention trials.</p>	
Funding Information:	<p>National Institute of Diabetes and Digestive and Kidney Diseases (U01 DK061010, U01 DK061034, U01 DK061042, U01 DK061058, U01 DK085465, U01 DK085453, U01 DK085461, U01 DK085466, U01 DK085499, U01 DK085504, U01 DK085509, U01 DK103180, U01 DK103153, U01 DK085476, U01 DK103266, U01 DK103282, U01 DK106984, U01 DK106994, U01 DK107013, U01 DK107014, UC4 DK106993)</p>	Not applicable
Author Comments:	No dualities of interest relevant to this article are reported by the Authors.	
Additional Information:		
Question	Response	
MANUSCRIPT HISTORY: Has this manuscript been submitted or reviewed before in an Endocrine Society journal?	No	
<p>WELLCOME TRUST / RESEARCH COUNCILS UK:</p> <p>In accordance with Wellcome Trust and Research Councils UK policies, the Endocrine Society will deposit to PubMed Central the final published article. For the full policy, see the Author Guidelines. Indicate below if the paper you are submitting has received funding from any of the organizations listed below:</p>	None of the above	
<p>CELL LINE AUTHENTICATION:</p> <p>I have read and understood the Cell Line Authentication policy and describe my submission as follows:</p>	Not applicable to my manuscript.	
STEROID HORMONE MEASUREMENT:	Not applicable to my manuscript.	

<p>I have read and understood the Steroid Hormone Measurement policy and describe my submission as follows:</p>	
<p>PRECIS:</p> <p>Please submit a brief description of your paper that will appear on the Table of Contents along with the title, should your paper be accepted. The description should be NO LONGER THAN 200 CHARACTERS and should serve to buttress the content of the title by simply stating what was done and what was concluded.</p>	<p>We studied implications of age and islet autoantibody type on progression from single to multiple autoantibodies in relatives and found that risk differs according to the primary autoantigen involved</p>
<p>INVITED SUBMISSION: Is this an invited submission?</p>	<p>No</p>
<p>SPECIAL REQUESTS:</p> <p>Enter specific comments or requests to the editors here.</p>	

Impact of age and antibody type on progression from single to multiple autoantibodies in type 1 diabetes relatives (jc.2017-00569).

Dear Editor,

In response to the comments of reviewers, please find hereafter our comments:

Reviewer #1:

In addition to the 994 relatives positive for a single autoantibody (GADA, IA-2A or IAA) with normal oral glucose tolerance at baseline:

State how many of the 151,458 relatives screened in TrialNet PTP between March 1st 2004 and March 31st 2015 were:

- 1) single autoantibody-positive with dysglycaemia at baseline;**
- 2) multiple autoantibody-positive with normoglycaemia and dysglycaemia, respectively.**

Response: We appreciate the reviewer's request for clarification related to how this cohort compares to those who were not included. Of all subjects who were confirmed as positive for a single autoantibody, 994 were identified as normoglycaemic (presented in this manuscript) and an additional 276 subjects had abnormal glucose tolerance. We have included these clarifications in the text of the paper (line 136-138).

We have specifically not included information related to those subjects who were positive for multiple autoantibodies; that cohort is being described through another manuscript.

State how many of the 994 relatives were at baseline ZnT8A positive and negative, respectively, and consider this status in the analysis and interpretation.

Response: As reported in the methods section (lines 102-104), participants included in this study were those screened and confirmed positive for only one autoantibody (GADA, IAA or IA-2A) and were also additionally tested for islet cell antibodies (ICA) and ZnT8A. Therefore, all 994 relatives included in these analyses were ZnT8A-negative, since any subjects who tested positive for ZnT8A (or any other additional autoantibody) were classified as multiple antibody-positive.

Clarify the primary outcome: "... confirmed development of multiple autoantibodies defined as detection on two (subsequent?) occasions ..."

Response: We again appreciate the request for clarification on this important point. Indeed, the primary endpoint for this paper is confirmed development of multiple autoantibodies, which we have defined as being identified as having two or more positive autoantibodies (GADA, IAA, IA-2A, ZnT8A, or ICA). This positivity is required to be confirmed on two consecutive evaluations that are done within one year of each other. The date the subject was identified as confirmed positive for multiple autoantibodies is the date of the first multiple antibody positive result that was subsequently confirmed. This has been clarified in the text (lines 120-121).

Importantly, the authors should include HLA in their analysis.

Response: Analyses of the associations between HLA, autoantibody type and risk of progression in this group of confirmed single antibody-positive relatives have already been published in Bingley PJ et al. Diabetologia 2016;59:542-549 (ref 16). We do not therefore feel it is appropriate to add it to the present report. We have however highlighted the previous analysis in the discussion (line 183-185).

Reviewer #2:

It would be helpful briefly to describe the technique of recursive partitioning analysis.

Response: We thank the reviewer for this helpful suggestion, and have included some clarification on this in the statistical methods section (lines 128-130). Overall, recursive partitioning is a model-based and iterative approach that can be used to identify cutpoints of a marker that best differentiate prognosis based on the outcome of interest, here, risk of progression to multiple positive autoantibodies.

In Figures 1 and 2 the decimal points on the y-axis are unnecessary.

Response: We acknowledge that decimal points are not necessarily required in these time-to-event Kaplan-Meier plots for the distributions of time to progression to multiple positive autoantibodies. However, their inclusion is not erroneous. Nonetheless, we modified the formatting of the y-axis for Figures 2a and 2b to make it more readable.

Perhaps add a rider that we do not know if this information necessarily holds true for older people developing type 1 diabetes in their thirties or forties.

Response: We acknowledge that many of these subjects included in our analysis are younger. However, the age range for subjects in these analyses is from 1 to 51 years old at positive autoantibody determination. In fact, 375 of these subjects were 30 years or older. While we did not conduct a subset analysis focusing on these older subjects, they are indeed well represented in this cohort.



1 **Full title**

2 Impact of age and antibody type on progression from single to multiple autoantibodies in type 1
3 diabetes relatives

4 **Short title**

5 Autoantibody progression in pre-type 1 diabetes

6 **Authors**

7 Emanuele Bosi¹, David C Boulware², Dorothy J Becker³, Jane H Buckner⁴, Susan Geyer², Peter A
8 Gottlieb⁵, Courtney Henderson², Amanda Kinderman², Jay M Sosenko⁶, Andrea K Steck⁵, Polly J
9 Bingley⁷; Type 1 Diabetes TrialNet Study Group*.

10 **Affiliations**

11 ¹ Diabetes Research Institute, San Raffaele Hospital and San Raffaele Vita Salute University, Milan,
12 Italy. bosi.emanuele@hsr.it

13 ² Division of Informatics and Biostatistics, University of South Florida, Tampa, FL, USA.

14 ³ Department of Pediatric Endocrinology, Children's Hospital of Pittsburgh, University of
15 Pittsburgh, Pittsburgh, PA, USA

16 ⁴ Translational Immunology Program, Benaroya Research Institute, Seattle, WA, USA

17 ⁵ Barbara Davis Center for Childhood Diabetes, University of Colorado School of Medicine, Aurora,
18 CO, USA

19 ⁶ Division of Endocrinology, University of Miami, Miami, FL, USA

20 ⁷ School of Clinical Sciences, University of Bristol, Bristol, UK.

21 * A complete list of the Type 1 Diabetes TrialNet Study Group can be found in the Supplementary
22 Data online

23

24 **Funding**

25 The sponsor of the trial was the Type 1 Diabetes TrialNet Study Group. Type 1 Diabetes TrialNet Study
26 Group is a clinical trials network funded by the National Institutes of Health (NIH) through the National
27 Institute of Diabetes and Digestive and Kidney Diseases, the National Institute of Allergy and Infectious
28 Diseases, and The Eunice Kennedy Shriver National Institute of Child Health and Human Development,
29 through the cooperative agreements U01 DK061010, U01 DK061034, U01 DK061042, U01 DK061058, U01
30 DK085465, U01 DK085453, U01 DK085461, U01 DK085466, U01 DK085499, U01 DK085504, U01 DK085509,
31 U01 DK103180, U01 DK103153, U01 DK085476, U01 DK103266, U01 DK103282, U01 DK106984, U01
32 DK106994, U01 DK107013, U01 DK107014, UC4 DK106993, and the Juvenile Diabetes Research Foundation
33 International (JDRF). The contents of this Article are solely the responsibility of the authors and do not
34 necessarily represent the official views of the NIH or the JDRF.

35

36 **Duality of interest**

37 No dualities of interest relevant to this article are reported by the Authors.

38

39 **Corresponding Author**

40 Emanuele Bosi, M.D.
41 Diabetes Research Institute
42 San Raffaele Hospital and Vita-Salute University
43 Via Olgettina, 60 - 20132 Milan, Italy
44 bosi.emanuele@hsr.it

45

46 **Word count: 2048 (main text)**

47 **Context.** Islet autoantibodies are markers of type 1 diabetes and an increase in number of
48 autoantibodies detected during the preclinical phase is predictive of progression to overt disease.

49 **Objective.** To refine the impact of age in relation to islet antibody type on the progression from
50 single to multiple autoantibodies in relatives of people with type 1 diabetes.

51 **Research design and methods.** We examined 994 relatives with normal glucose tolerance and
52 positive for a single autoantibody, followed prospectively in the TrialNet Pathway to Prevention.
53 Antibodies to GAD (GADA), insulin (IAA), IA-2 (IA-2A), zinc transporter 8 (ZnT8A) and ICA were
54 tested every 6-12 months. The primary outcome was confirmed development of multiple
55 autoantibodies. Age was categorized as <8yr; 8-11yr; 12-17yr; ≥18yr and optimal age breakpoints
56 identified by recursive partitioning analysis.

57 **Results.** After median follow-up of 2 years, 141 relatives had developed ≥ 1 additional
58 autoantibodies. Five-year risk was inversely related to age, but the pattern differed by antibody
59 type: relatives with GADA showed a gradual decrease in risk over the four age groups, while
60 relatives with IAA showed a sharp decrease above the age of 8 years. Recursive partitioning
61 analysis identified age breakpoints at 14 years in relatives with GADA and at 4 years in relatives
62 with IAA.

63 **Conclusions.** In relatives with IAA, spread of islet autoimmunity is largely limited to early
64 childhood, while immune responses initially directed at GAD can mature over a longer period of
65 time. These differences have important implications for monitoring these subjects and for
66 designing prevention trials.

67 **Introduction**

68 Islet autoimmunity leading to type 1 diabetes develops and progresses silently over many years
69 before glucose intolerance and symptomatic hyperglycemia occur (1). Islet autoantibodies are the
70 best validated markers of this ongoing pathogenetic process and are used to predict clinical
71 disease (2) and stage its preclinical phase (3).

72 Several prospective studies, including those following infants at genetic risk from birth, have
73 shown that the antibody response against β -cells within pancreatic islets usually targets several
74 autoantigens including insulin (IAA), glutamic acid decarboxylase (GADA), insulinoma-associated
75 antigen 2/ICA512 (IA2A) and zinc transporter 8 (ZnT8A), in varying sequence. Maturation of this
76 humoral immune response, as shown by increasing autoantibody number, titer and affinity, is
77 associated with an increased risk of progression to the disease (4).

78 Specifically, the number of autoantibodies detected seems crucial for the prediction of disease,
79 with a relatively low risk associated with positivity for a single autoantibody, increasing to near
80 certainty of developing type 1 diabetes following the appearance of multiple (i.e. two or more)
81 autoantibodies (4-13). Therefore, seroconversion from single to multiple autoantibodies appears
82 to be the hallmark of a 'point of no return' in the type 1 diabetes pathogenetic process, marking
83 the transition from a state of predisposition to a preclinical stage of the disease (3). Associations
84 have been described with younger age and HLA class II genotype (6, 10), but the determinants of
85 this maturation remain largely unknown.

86 Previous analyses in relatives followed prospectively in the TrialNet Pathway to Prevention (PTP)
87 (formerly Natural History Study) (14,15) have indicated heterogeneity in the development and
88 spreading of islet autoantibody responses, and have also shown that progression from single to
89 multiple autoantibodies is not restricted to early childhood and high risk genotypes (16,17). In light
90 of this evidence and the implications for design of targeted interventions to prevent or delay
91 progression of islet autoimmunity, we have undertaken a more in-depth investigation into the
92 transition from single to multiple autoantibodies, with specific regard to the impact of age in
93 relation to antibody type.

94 **Methods**

95 **Study population**

96

97 Non-diabetic first, second and third degree relatives of people with type 1 diabetes were recruited
98 to the TrialNet PTP (ClinicalTrials.gov identifier: NCT00097292) as previously described (14). All
99 study participants gave informed consent and the Ethics Committee responsible for each clinical
100 site approved the study. Participants were included in this analysis if they had antibodies to the
101 same single islet autoantigen (GADA, IAA or IA-2A) detected on at least two occasions, and
102 antibody results were available from at least one subsequent study visit. All samples were
103 screened for GADA, IAA and IA-2A, and if levels of any of these were above the threshold of
104 positivity, samples were additionally tested for islet cell antibodies (ICA) and ZnT8A. Individuals

105 with confirmed islet autoantibodies underwent baseline assessment including oral glucose
106 tolerance testing and were followed every 6–12 months in accordance with the PTP study
107 protocol, as previously reported (14). Relatives with the protective HLA DQB1*0602 allele were
108 not included in this analysis.

109

110 Assays

111

112 GADA, IAA, IA-2A and ZnT8A were measured by radioimmunoassay in the TrialNet Core laboratory
113 at the Barbara Davis Center for Childhood Diabetes (BDC), Denver, CO and ICA by indirect
114 immunofluorescence at the University of Florida, Gainesville, FL, as previously described (18-20).

115

116 Statistical analysis

117

118 The primary outcome of the analysis was confirmed development of multiple autoantibodies
119 defined as detection on two occasions of at least two of the five islet autoantibodies included in
120 the testing strategy (GADA, IAA, IA-2A, ZnT8A and ICA); this confirmation is based on two
121 consecutive autoantibody tests that are done within one year of each other. The time-to-event
122 was calculated from the date of first detection of a single islet autoantibody to date of first
123 detection of multiple autoantibodies. The risk of developing multiple autoantibodies was assessed
124 by survival analysis using Kaplan–Meier curves. Cox proportional hazards regression models were
125 used for multivariable analysis. Age groups were initially categorized using boundaries based on
126 common definitions of infancy to pre-pubertal, adolescence and adulthood (<8yr; 8-11yr; 12-17yr;
127 ≥18yr); and then refined by recursive partitioning analysis used to identify optimal age breakpoints
128 within each single autoantibody population (21-23). Recursive partitioning is a model-based
129 method used to identify a cutpoint for a marker that best differentiates subjects in relation to an
130 outcome of interest, such as risk of progression to multiple positive autoantibodies.

131

132 Results

133

134 Of 151,458 relatives screened in the TrialNet PTP between March 1st 2004 and March 31st 2015,
135 994 were positive for a single autoantibody (GADA, IAA or IA-2A) with normal oral glucose
136 tolerance at baseline and were therefore eligible for inclusion in the analysis, while an additional
137 276 subjects had abnormal glucose tolerance. Of the 994 positive for a single autoantibody with
138 normal glucose tolerance, 709 (71.3%) had GADA, 236 (23.7%) had IAA and 49 (4.9%) had IA-2A;
139 59.6% were female.

140

141 The median age of the participants was 17.6 years (interquartile range 9.8 - 36.2); 183 (18.4%)
142 were aged less than 8 years, 157 (15.8%) between 8 and 11 years, 169 (17.0%) between 12 and 17
143 years and 485 (48.8%) 18 years or more.

144

145 After a median follow-up of 2.0 years (IQR 0.8-3.8), 141 relatives had developed ≥ 1 additional
146 autoantibodies. Estimated cumulative risk within 5 years was 23% (95% CI 19-27) overall, and did

not vary between autoantibody types (GADA 25% [CI 20-30]; IAA 19% [CI 11-27]; IA-2A 23% [CI 7-38]; $p=0.09$).

The overall risk of developing additional autoantibodies was inversely related to age (multivariable HR 0.96, 95% CI 0.94-0.99, $p=0.005$), but heterogeneity between autoantibody type was identified. Table 1 shows the risk of developing multiple autoantibodies within 5 years categorized by age. Relatives with GADA showed a gradual decrease in risk across the four age groups, while relatives with IAA showed a sharp decrease in risk above the age of 8 years. Analysis between age groups was not performed in relatives with IA-2A due to an insufficient number of subjects.

The age distribution of the relatives who developed multiple autoantibodies also differed between those with GADA or IAA (Figure 1). Relatives below 8 years of age represented 71% of the IAA-positive individuals who became multiple autoantibody-positive within 5 years, while the age of GADA-positive individuals who progressed were more evenly distributed throughout childhood and adolescence. Of the relatives who developed one or more additional autoantibodies, 17/24 (71%) with IAA vs 22/99 (22%) with GADA were aged <8 years ($p=0.048$).

Recursive partitioning analysis confirmed differences in age-related risk between GADA-positive and IAA-positive groups and identified age breakpoints at 14 years in relatives with GADA alone and 4 years in those with IAA alone. The survival time from initial detection of a single islet autoantibody to first detection of multiple antibodies in subgroups categorized by these age breakpoints for GADA-positive relatives is shown in Figure 2A, and for IAA-positive relatives in Figure 2B. Among the GADA-positive relatives, 99 individuals developed additional autoantibodies within 5 years, of whom 59 were aged less than 14 years and 40 were aged ≥ 14 years (estimated cumulative risk 35.2% [95% CI] (27.9, 43.8) vs 17.7% [95% CI] (12.6, 24.6), respectively; logrank test $p<0.001$). Among IAA-positive relatives, 24 individuals developed additional autoantibodies within 5 years, of whom 12 were aged less than 4 years and 12 were aged ≥ 4 years (estimated cumulative risk 73.1% [95% CI] (48.5, 92.6) vs 11.4% [95% CI] (6.3, 20.4), respectively; logrank test, $p<0.001$).

Discussion

The main finding of this analysis is that the interaction between age and risk of progression from single to multiple autoantibodies differs according to whether the primary autoantigen is GAD or insulin. Specific age breakpoints were very different in those initially positive for IAA compared to those with GADA. This suggests that early interventions targeting single autoantibody-positive individuals at risk for type 1 diabetes may differ in their effectiveness at different ages, depending on whether they have GADA or IAA and provides some guidance as to what age groups to target.

Previous analyses in relatives followed prospectively in the TrialNet PTP (14,15) found that, in those positive for a single autoantibody, the risk of progression to multiple antibodies was 22% within 5 years and to type 1 diabetes was 6% (16). As confirmed in the present analysis, overall risk of development of multiple autoantibodies was independent of antibody type, inversely related to age, and associated with high and intermediate risk HLA Class II genotypes and high

186 GADA titers (16). In addition, progression appeared not to be influenced by initial body mass
187 index or other metabolic variables (24). More recently, recursive partitioning analysis
188 demonstrated that age and GADA titer taken together were helpful in stratifying the overall risk of
189 progression from single to multiple autoantibodies (17).

190 These additional analyses focus on evaluation of the time course of progression among single
191 autoantibody-positive subjects, depending whether the first autoantibodies detected were GADA
192 or IAA. These demonstrate marked differences between the initial IAA and GADA groups. Although
193 risk was inversely related to age in both groups, the relationships were not the same; while risk in
194 GADA-positive relatives decreased gradually up to age 18 years, in IAA-positive relatives risk was
195 concentrated in younger children with high risk of progression up to age 8 years, and a sharp
196 decline thereafter. This pattern was confirmed by recursive partitioning analysis which showed
197 significant differences in risk and identified different age breakpoints for IAA (4 years) and for
198 GADA (14 years).

199 Based on these observations, spread of autoimmunity from insulin to other islet antigens seems
200 largely limited to early childhood, while autoimmune responses initially directed against GAD can
201 mature over a longer period of time, including the whole of adolescence. It will be interesting to
202 see whether this pattern is confirmed in studies of cellular immune responses.

203 These findings are relevant to the design of prevention trials targeting the early phases of type 1
204 diabetes-associated autoimmunity. In individuals with IAA alone, intervention should be
205 considered only below the age of 8 years, with the highest priority given to individuals below 4
206 years of age; conversely, in individuals with GADA alone, intervention seems justified up to the age
207 of 18 years, with the highest potential below 14 years. Although there is no sharp decline after age
208 18, the overall risk of developing additional antibodies is relatively low in GADA-positive adults
209 (16% within 5 years) and we suggest that issues of power and the sample size required should be
210 carefully evaluated before including them in type 1 diabetes prevention trials.

211 A particular strength of this study is the size of the cohort, screened and followed up every 6-12
212 months according to a standard protocol (14). One limitation is that, in contrast with studies
213 following infants from birth, the time at which seroconversion has occurred is unknown when
214 relatives are found to be positive for an autoantibody in the TrialNet PTP. This may potentially lead
215 to underestimation of the duration of single autoantibody positivity but, since the focus of this
216 study is the time to further seroconversion to multiple autoantibodies, the time of initial
217 seroconversion is less relevant in this context. As in other TrialNet studies, we have included
218 appearance of ICA in the definition of development of multiple autoantibodies because, although
219 GADA and I-2A contribute to ICA staining (25,26), there is not complete overlap and previous
220 analyses in the PTP cohort have demonstrated that ICA are associated with additional risk (20).

221 Although the determinants of islet autoimmunity underlying type 1 diabetes remain unknown,
222 accumulating evidence suggests that it develops through multistep involvement of several
223 pathogenetic pathways (27). Accordingly, interventions at an earlier stage might rely on simpler
224 approaches, such as antigen based therapies, while at a later stage, a more complex approach,

225 based for instance on combination therapies, is likely to be necessary to halt or delay the
226 progression of the disease process (28). Based on our current ability to predict type 1 diabetes,
227 single autoantibody positivity is the earliest detectable sign of the ongoing autoimmune process,
228 when the chances of a successful intervention could be greatest. The evidence of heterogeneity in
229 the progression of islet autoimmunity in relatives of the TrialNet PTP study, the largest study
230 cohort ever screened and followed up, calls for different therapeutic approaches in single
231 autoantibody positive individuals with IAA or GADA, with the highest potential for success in
232 interventions designed for early childhood in IAA-positive and no later than adolescence in GADA-
233 positive individuals.

234 **Author contribution**

235 All authors were members of the TrialNet Study Group and contributed to the data used in this
236 article. EB and PJB wrote the manuscript. PJB and DCB designed and conducted the statistical
237 analysis. All authors contributed to discussion, reviewed/edited the manuscript and gave final
238 approval for the paper to be published. SG is the guarantor of this work and, as such, had full
239 access to all the data in the study and takes responsibility for the integrity of the data and the
240 accuracy of the data analysis.

241 **References**

- 242 1) Atkinson MA, Eisenbarth GS, Michels AW. Type 1 diabetes. *Lancet* 2014;383:69-82
- 243 2) Bonifacio E. Predicting type 1 diabetes using biomarkers. *Diabetes Care* 2015;38:989-996
- 244 3) Insel RA, Dunne JL, Atkinson MA, Chiang JL, Dabelea D, Gottlieb PA, Greenbaum CJ, Herold
245 KC, Krischer JP, Lernmark Å, Ratner RE, Rewers MJ, Schatz DA, Skyler JS, Sosenko JM,
246 Ziegler AG. Staging presymptomatic type 1 diabetes: a scientific statement of JDRF, the
247 Endocrine Society, and the American Diabetes Association. *Diabetes Care* 2015;38:1964-
248 1974
- 249 4) Ziegler AG, Rewers M, Simell O, Simell T, Lempainen J, Steck A, Winkler C, Ilonen J, Veijola
250 R, Knip M, Bonifacio E, Eisenbarth GS. Seroconversion to multiple islet autoantibodies and
251 risk of progression to diabetes in children. *JAMA* 2013;309:2473-2479
- 252 5) Orban T, Sosenko JM, Cuthbertson D, Krischer JP, Skyler JS, Jackson R, Yu L, Palmer JP,
253 Schatz D, Eisenbarth G; Diabetes Prevention Trial-Type 1 Study Group. Pancreatic islet
254 autoantibodies as predictors of type 1 diabetes in the Diabetes Prevention Trial-Type 1.
255 *Diabetes Care* 2009;32:2269-2274
- 256 6) Ziegler AG, Bonifacio E; BABYDIAB-BABYDIET Study Group. Age-related islet autoantibody
257 incidence in offspring of patients with type 1 diabetes. *Diabetologia* 2012;55:1937-1943
- 258 7) Steck AK, Johnson K, Barriga KJ, Miao D, Yu L, Hutton JC, Eisenbarth GS, Rewers MJ. Age of
259 islet autoantibody appearance and mean levels of insulin, but not GAD or IA-2
260 autoantibodies, predict age of diagnosis of type 1 diabetes: diabetes autoimmunity study in
261 the young. *Diabetes Care* 2011;34:1397-1399
- 262 8) Parikka V, Näntö-Salonen K, Saarinen M, Simell T, Ilonen J, Hyöty H, Veijola R, Knip M,
263 Simell O. Early seroconversion and rapidly increasing autoantibody concentrations predict

- prepubertal manifestation of type 1 diabetes in children at genetic risk. *Diabetologia* 2012;55:1926-1936
- 9) Steck AK, Vehik K, Bonifacio E, Lernmark A, Ziegler AG, Hagopian WA, She J, Simell O, Akolkar B, Krischer J, Schatz D, Rewers MJ; TEDDY Study Group. Predictors of Progression From the Appearance of Islet Autoantibodies to Early Childhood Diabetes: The Environmental Determinants of Diabetes in the Young (TEDDY). *Diabetes Care* 2015;38:808-813
- 10) Krischer JP, Lynch KF, Schatz DA, Ilonen J, Lernmark Å, Hagopian WA, Rewers MJ, She JX, Simell OG, Toppari J, Ziegler AG, Akolkar B, Bonifacio E; TEDDY Study Group. The 6 year incidence of diabetes-associated autoantibodies in genetically at-risk children: the TEDDY study. *Diabetologia* 2015;58:980-987
- 11) Achenbach P, Hummel M, Thümer L, Boerschmann H, Höfelmann D, Ziegler AG. Characteristics of rapid vs slow progression to type 1 diabetes in multiple islet autoantibody-positive children. *Diabetologia*. 2013;56:1615-1622
- 12) Chmiel R, Giannopoulou EZ, Winkler C, Achenbach P, Ziegler AG, Bonifacio E. Progression from single to multiple islet autoantibodies often occurs soon after seroconversion: implications for early screening. *Diabetologia*. 2015;58:411-3
- 13) Steck AK, Dong F, Waugh K, Frohnert BI, Yu L, Norris JM, Rewers MJ. Predictors of slow progression to diabetes in children with multiple islet autoantibodies. *J Autoimmun*. 2016;72:113-117
- 14) Mahon JL, Sosenko JM, Rafkin-Mervis L, Krause-Steinrauf H, Lachin JM, Thompson C, Bingley PJ, Bonifacio E, Palmer JP, Eisenbarth GS, Wolfsdorf J, Skyler JS; TrialNet Natural History Committee; Type 1 Diabetes TrialNet Study Group. The TrialNet Natural History Study of the Development of Type 1 Diabetes: objectives, design, and initial results. *Pediatr Diabetes* 2009;10:97-104
- 15) Vehik K, Beam CA, Mahon JL, Schatz DA, Haller MJ, Sosenko JM, Skyler JS, Krischer JP; TrialNet Natural History Study Group. Development of autoantibodies in the TrialNet Natural History Study. *Diabetes Care*. 2011;34:1897-1901
- 16) Bingley PJ, Boulware DC, Krischer JP; Type 1 Diabetes TrialNet Study Group. The implications of autoantibodies to a single islet antigen in relatives with normal glucose tolerance: development of other autoantibodies and progression to type 1 diabetes. *Diabetologia* 2016;59:542-549
- 17) Xu P, Krischer JP. Prognostic Classification Factors Associated With Development of Multiple Autoantibodies, Dysglycemia, and Type 1 Diabetes-A Recursive Partitioning Analysis. *Diabetes Care* 2016;39:1036-1044
- 18) Yu L, Rewers M, Gianani R, Kawasaki E, Zhang Y, Verge C, Chase P, Klingensmith G, Erlich H, Norris J, Eisenbarth GS. Antiislet autoantibodies develop sequentially rather than simultaneously. *J Clin Endocrinol Metab* 1996;81:4264-4267
- 19) Bonifacio E, Yu L, Williams AK, Eisenbarth GS, Bingley PJ, Marcovina SM, Adler K, Ziegler AG, Mueller PW, Schatz DA, Krischer JP, Steffes MW, Akolkar B. Harmonization of glutamic acid decarboxylase and islet antigen-2 autoantibody assays for national institute of

- diabetes and digestive and kidney diseases consortia. *J Clin Endocrinol Metab.* 2010;95:3360-3367
- 20)** Yu L, Boulware DC, Beam CA, Hutton JC, Wenzlau JM, Greenbaum CJ, Bingley PJ, Krischer JP, Sosenko JM, Skyler JS, Eisenbarth GS, Mahon JL; Type 1 Diabetes TrialNet Study Group. Zinc transporter-8 autoantibodies improve prediction of type 1 diabetes in relatives positive for the standard biochemical autoantibodies. *Diabetes Care* 2012;35:1213–1218
- 21)** Ciampi IA, Thiffault J. Recursive partition and amalgamation (RECPAM) for censored survival data: criteria for tree selection. *Stat Softw Newsletter* 1988;14:78–81
- 22)** Gordon L, Olshen RA. Tree-structured survival analysis. *Cancer Treat Rep* 1985;69:1065–1069
- 23)** Breiman L, Friedman JH, Olshen RA, Stone CJ. *Classification and Regression Trees*. Belmont, CA, Wadsworth International Group, 1984
- 24)** Meah FA, DiMeglio LA, Greenbaum CJ, Blum JS, Sosenko JM, Pugliese A, Geyer S, Xu P, Evans-Molina C; Type 1 Diabetes TrialNet Study Group. The relationship between BMI and insulin resistance and progression from single to multiple autoantibody positivity and type 1 diabetes among TrialNet Pathway to Prevention participants. *Diabetologia* 2016;59:1186-1195
- 25)** Richter W, Seissler J, Northemann W, Wolfahrt S, Meinck HM, Scherbaum WA. Cytoplasmic islet cell antibodies recognize distinct islet antigens in IDDM but not in stiff man syndrome. *Diabetes.* 1993;42:1642-1648.
- 26)** Bonifacio E, Lampasona V, Genovese S, Ferrari M, Bosi E. Identification of protein tyrosine phosphatase-like IA2 (islet cell antigen 512) as the insulin-dependent diabetes-related 37/40K autoantigen and a target of islet-cell antibodies. *J Immunol.* 1995;155:5419-1526
- 27)** Bluestone JA, Herold K, Eisenbarth G. Genetics, pathogenesis and clinical interventions in type 1 diabetes. *Nature* 2010;464:1293-1300
- 28)** Skyler JS. Prevention and reversal of type 1 diabetes--past challenges and future opportunities. *Diabetes Care* 2015;38:997-1007

334 **Legend to table and figures**

335

336 **Table 1.** Risk of developing multiple autoantibodies within 5 yr in single autoantibody-positive
337 relatives categorized by age (95% CI).

338

339 **Figure 1:** GADA-positive (panel A, n=99) and IAA-positive participants (panel B, n=24) who
340 developed additional autoantibodies within 5 years subdivided by age group. Among the relatives
341 who progressed, 17/24 (71%) with IAA vs. 27/99 (27%) GADA were aged <8 years ($p<0.05$).

342

343 **Figure 2.-** Time from initial detection of: **A)** single GADA to first detection of multiple antibodies in
344 relatives aged <14 years (black line) and ≥ 14 years (grey line); and **B)** of single IAA to first detection
345 of multiple antibodies in relatives aged <4 years (black line) and ≥ 4 years (grey line)

346

347 **Table 1** - Risk of developing multiple autoantibodies within 5 yr in single autoantibody-positive
348 relatives categorized by age (95% CI)

349

<u>Antibody type</u>	Age at entry (n)				Overall
	<8 yr	8-11 yr	12-17yr	>18 yr	
GADA (n)	107	113	117	372	709
- Risk	35% (24-46)	38% (24-51)	28% (14-41)	16% (10-22)	25% (20-30)
IAA (n)	69	33	43	91	236
- Risk	37% (22-52)	4% (0-11)	13% (0-34)	13% (0-25)	19% (11-27)
IA-2A (n)	7	11	9	22	49
- Risk	<i>Insufficient number of subjects to assess</i>				23% (7-38)


350

351

Table 1 - Risk of developing multiple Ab within 5 yr in single Ab-positive relatives categorized by age (95% CI)

Ab type	Age at entry (n)				Overall
	<8 yr	8-11 yr	12-17yr	>18 yr	
GADA (n)	107	113	117	372	709
- Risk	35% (24-46)	38% (24-51)	28% (14-41)	16% (10-22)	25% (20-30)
IAA (n)	69	33	43	91	236
- Risk	37% (22-52)	4% (0-11)	13% (0-34)	13% (0-25)	19% (11-27)
IA-2A (n)	7	11	9	22	49
- Risk	<i>Insufficient number of subjects to assess</i>				23% (7-38)

A



B

